

*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name:	NEBuffer™ r2.1
Catalog Number:	B6002S
Concentration:	10 X Concentrate
Packaging Lot Number:	10150648
Expiration Date:	03/2025
Storage Temperature:	-20°C
Specification Version:	PS-B6002S v1.0
Composition (1X):	10 mM Tris-HCl, 50 mM NaCl, 10 mM MgCl2, 100 μg/ml rAlbumin, (pH 7.9 @ 25°C)

NEBuffer™ r2.1 Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
B6002SVIAL	NEBuffer™ r2.1	10138406	Pass	

Assay Name/Specification	Lot # 10150648
Conductivity (buffers/solutions)	Pass
The conductivity of 10X NEBuffer™ r2.1 is between 55 and 62 mS/cm at 25°C.	
Functional Testing (Restriction Digest, Buffer)	Pass
A 50 µl reaction in 1X NEBuffer™ r2.1 containing 1 µg of Lambda DNA and 1 unit of	
SphI incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.	
Functional Testing (Restriction Digest, Buffer)	Pass
A 50 µl reaction in 1X NEBuffer™ r2.1 containing 1 µg of Lambda DNA and 1 unit of HindIII incubated for 1 hour at 37°C results in complete digestion of the substrate	
DNA as determined by agarose gel electrophoresis.	
pH (buffers/solutions)	Pass
The pH of 10X NEBuffer™ r2.1 is between pH 7.8 and 8.0 at 25°C.	
Non-Specific DNase Activity (16 hour, Buffer)	Pass
A 50 µl reaction in 1X NEBuffer™ r2.1 containing 1 µg of PhiX174-HaeIII DNA	1 400
incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease	
degradation as determined by agarose gel electrophoresis.	
RNase Activity (Buffer)	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10150648
A 10 µl reaction in 1X NEBuffer™ r2.1 containing 40 ng of a 300 base single-stranded RNA is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	
Endonuclease Activity (Nicking, Buffer) A 50 µl reaction in 1X NEBuffer™ r2.1 containing 1 µg of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Bu Rear

Ben Penta Production Scientist 11 May 2022

Erin Varney

Packaging Quality Control Inspector 11 May 2022

