

New England Biolabs Certificate of Analysis

Product Name: Shrimp Alkaline Phosphatase (rSAP)
Catalog Number: M0371S
Concentration: 1,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme that hydrolyzes 1 μmol of p-Nitrophenyl Phosphate, PNPP in a total reaction volume of 1 ml in 1 minute at 37°C
Packaging Lot Number: 10153699
Expiration Date: 12/2023
Storage Temperature: -20°C
Storage Conditions: 25 mM Tris-HCl, 1 mM MgCl₂, 50 % Glycerol, (pH 7.5 @ 25°C)
Specification Version: PS-M0371S/L v1.0


Shrimp Alkaline Phosphatase (rSAP) Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0371SVIAL	Shrimp Alkaline Phosphatase (rSAP)	10124428	Pass
B6004SVIAL	rCutSmart™ Buffer	10149689	Pass

Assay Name/Specification	Lot # 10153699
<p>Non-Specific DNase Activity (16 Hour) A 50 μl reaction in NEBuffer 4 containing 1 μg of PhiX174-HaeIII DNA and a minimum of 10 units of Shrimp Alkaline Phosphatase (rSAP) incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass
<p>Exonuclease Activity (Radioactivity Release) A 50 μl reaction in CutSmart® Buffer containing 1 μg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 10 units of Shrimp Alkaline Phosphatase (rSAP) incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.</p>	Pass
<p>Protein Purity Assay (SDS-PAGE) Shrimp Alkaline Phosphatase (rSAP) is \geq 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass
<p>RNase Activity (Extended Digestion) A 10 μl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 μl of Shrimp Alkaline Phosphatase (rSAP) is incubated at 37°C.</p>	Pass


Assay Name/Specification	Lot # 10153699
<p>After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p> <p>Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart[®] Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 5 units of Shrimp Alkaline Phosphatase (rSAP) incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	<p>Pass</p>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Ana Egana
Production Scientist
31 May 2022



Erin Varney
Packaging Quality Control Inspector
31 May 2022