

## New England Biolabs Certificate of Analysis

**Product Name:** LunaScript® RT SuperMix  
**Catalog Number:** M3010X  
**Concentration:** 5 X Concentrate  
**Packaging Lot Number:** 10178622  
**Expiration Date:** 06/2024  
**Storage Temperature:** -20°C  
**Specification Version:** PS-M3010S/L/X/E v2.0  
**Composition (1X):** Proprietary

LunaScript® RT SuperMix Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M3010LVIAL	LunaScript® RT SuperMix	10151266	Pass

Assay Name/Specification	Lot # 10178622
<b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in ThermoPol® Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 100 units of Luna® Reverse Transcriptase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	<b>Pass</b>
<b>Protein Purity Assay (SDS-PAGE)</b> Luna® Reverse Transcriptase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	<b>Pass</b>
<b>Phosphatase Activity (pNPP)</b> A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl <sub>2</sub> containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 100 units of Luna® Reverse Transcriptase incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	<b>Pass</b>
<b>RNase Activity Assay (4 Hour Digestion)</b> A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 100 units of Luna® Reverse Transcriptase is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	<b>Pass</b>
<b>Single Stranded DNase Activity (FAM-Labeled Oligo)</b>	<b>Pass</b>

Assay Name/Specification	Lot # 10178622
<p>A 50 µl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent internal labeled oligonucleotide and a minimum of 250 units of Luna® Reverse Transcriptase incubated for 16 hours at 37°C yields &lt;5% degradation as determined by capillary electrophoresis.</p>	
<p><b>qPCR DNA Contamination (E. coli Genomic)</b> A minimum of 1 µl of LunaScript® RT SuperMix is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	<b>Pass</b>
<p><b>Functional Testing (Two-Step RT-qPCR)</b> The LunaScript® RT SuperMix is functionally tested in two-step RT-qPCR with human RNA template, resulting in a standard curve with a calculated qPCR efficiency of 90-110%, and a dynamic range of 7 orders of magnitude.</p>	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Trinh Nguyen  
Production Scientist  
28 Jun 2022



Michael Tonello  
Packaging Quality Control Inspector  
30 Jan 2023