

New England Biolabs Certificate of Analysis

Product Name: Deoxynucleotide (dNTP) Solution Mix
Catalog Number: N0447S
Concentration: 10 mM
Unit Definition: N/A
Packaging Lot Number: 10159216
Expiration Date: 03/2024
Storage Temperature: -20°C
Storage Conditions: Supplied in Ultrapure water as a sodium salt (pH 7.5)
Specification Version: PS-N0447S/L v4.0

Deoxynucleotide (dNTP) Solution Mix Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
N0447SVIAL	Deoxynucleotide (dNTP) Solution Mix	10147823	Pass

Assay Name/Specification	Lot # 10159216
PCR Amplification (2.0 kb Lambda, dNTPs) A 50 µl reaction in ThermoPol® Reaction Buffer in the presence of 200 µM Deoxynucleotide (dNTP) Solution Mix and 0.2 µM primers containing 1 ng Lambda DNA with 1.25 units of Taq DNA Polymerase for 25 cycles of PCR amplification results in the expected 2.0 kb product.	Pass
PCR Amplification (0.5 kb Lambda, dNTPs) A 50 µl reaction in ThermoPol® Reaction Buffer in the presence of 200 µM Deoxynucleotide (dNTP) Solution Mix and 0.2 µM primers containing 1 ng Lambda DNA with 1.25 units of Taq DNA Polymerase for 25 cycles of PCR amplification results in the expected 0.5 kb product.	Pass
PCR Amplification (5.0 kb Lambda, dNTPs) A 50 µl reaction in ThermoPol® Reaction Buffer in the presence of 200 µM Deoxynucleotide (dNTP) Solution Mix and 0.2 µM primers containing 1 ng Lambda DNA with 1.25 units of Taq DNA Polymerase for 25 cycles of PCR amplification results in the expected 5.0 kb product.	Pass
qPCR DNA Contamination (E. coli Genomic) A minimum of 1 µl of Deoxynucleotide (dNTP) Solution Mix is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from	Pass

Assay Name/Specification	Lot # 10159216
<p>purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	
<p>Non-Specific DNase Activity (16 Hour) A 50 μl reaction in NEBuffer 2 containing 1 μg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 10 μl of Deoxynucleotide (dNTP) Solution Mix incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass
<p>RNase Activity (Extended Digestion) A 10 μl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 μl of Deoxynucleotide (dNTP) Solution Mix is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass
<p>Phosphatase Activity (pNPP) A 200 μl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl₂ containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 40 μl Deoxynucleotide (dNTP) Solution Mix incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.</p>	Pass
<p>Physical Purity (HPLC) Deoxynucleotide (dNTP) Solution Mix is $\geq 99\%$ pure as determined by HPLC analysis.</p>	Pass
<p>Endonuclease Activity (Nicking) A 50 μl reaction in NEBuffer 2 containing 1 μg of supercoiled PhiX174 DNA and a minimum of 10 μl of Deoxynucleotide (dNTP) Solution Mix incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Christie Vazquez

Christie Vazquez
Production Scientist
08 Sep 2022

Erin Varney

Erin Varney
Packaging Quality Control Inspector
08 Sep 2022