

New England Biolabs Certificate of Analysis

Product Name: PNGase A
Catalog Number: P0707L
Concentration: 5,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to remove > 95% of the carbohydrate from 1 µg of denatured recombinant Avidin produced in Maize in 1 hour at 37°C in a total reaction volume of 10 µl.
Packaging Lot Number: 10160069
Expiration Date: 07/2023
Storage Temperature: 4°C
Storage Conditions: 50 mM NaCl, 20 mM Tris-HCl, 5 mM EDTA, (pH 7.5 @ 25°C)
Specification Version: PS-P0707S/L v1.0

PNGase A Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
P0707LVIAL	PNGase A	10153978	Pass
B2704SVIAL	NP-40	10129282	Pass
B1720SVIAL	10X Glycobuffer 3	10120516	Pass
B1704SVIAL	Glycoprotein Denaturing Buffer	10153869	Pass

Assay Name/Specification	Lot # 10160069
Glycosidase Activity (β-N-Acetylgalactosaminidase) A 10 µl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled β-N-Acetylgalactosaminidase substrate (GalNAcβ1-4Galβ1-4Glc-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β-N-Acetylglucosaminidase) A 10 µl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled β-N-Acetylglucosaminidase substrate (GlcNAcβ1-4GlcNAcβ1-4GlcNAc-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β-Mannosidase) A 10 µl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled β-Mannosidase substrate (Manβ1-4Manβ1-4Man-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer	Pass

Assay Name/Specification	Lot # 10160069
chromatography.	
<p>Glycosidase Activity (β1-3 Galactosidase) A 10 μl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-3GlcNAcβ1-4Galβ1-4Glc-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β1-4 Galactosidase) A 10 μl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-4GlcNAcβ1-3Galβ1-4Glc-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α-Neuraminidase) A 10 μl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α-Neuraminidase substrate (Neu5Acα2-3Galβ1-3GlcNAcβ1-3Galβ1-4Glc-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α-N-Acetylgalactosaminidase) A 10 μl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α-N-Acetylgalactosaminidase substrate (GalNAcα1-3(Fucaα1-2)Galβ1-4Glc-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β-Xylosidase) A 10 μl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled β-Xylosidase substrate (Xylβ1-4Xylβ1-4Xylβ1-4Xyl-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-2 Fucosidase) A 10 μl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α-Fucosidase substrate (Fucaα1-2Galβ1-4Glc-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α-Glucosidase) A 10 μl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α-Glucosidase substrate (Glcα1-6Glcα1-4Glc-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass


Assay Name/Specification	Lot # 10160069
<p>Glycosidase Activity (α1-6 Mannosidase) A 10 μl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α-Mannosidase substrate (Manα1-6Manα1-6(Manα1-3)Man-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-6 Galactosidase) A 10 μl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α-Galactosidase substrate (Galα1-6Galα1-6Glcα1-2Fru-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-3 Mannosidase) A 10 μl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α-Mannosidase substrate (Manα1-3Manβ1-4GlcNAc-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-3 Fucosidase) A 10 μl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α-Fucosidase substrate (Fuα1-3Galβ1-4GlcNAcβ1-3Galβ1-4Glc-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-3 Galactosidase) A 10 μl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α-Galactosidase substrate (Galα1-3Galβ1-4GlcNAc-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Protease Activity (SDS-PAGE) A 20 μl reaction in 1X Glyco Buffer 3 containing 24 μg of a standard mixture of proteins and a minimum of 25 units of PNGase A incubated for 20 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection.</p>	Pass
<p>Protein Purity Assay (SDS-PAGE) PNGase A is \geq 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Alicia Bielik
Production Scientist
02 Sep 2022



Erin Varney
Packaging Quality Control Inspector
02 Sep 2022