

New England Biolabs Certificate of Analysis

Product Name: Proteinase K, Molecular Biology Grade
Catalog Number: P8107S
Concentration: 800 U/ml
Unit Definition: One unit will digest urea-denatured hemoglobin at 37°C (pH 7.5) per minute to produce equal absorbance as 1.0 µmol L-tyrosine using Folin & Ciocalteu's phenol reagent.
Packaging Lot Number: 10156147
Expiration Date: 05/2025
Storage Temperature: -20°C
Storage Conditions: 20 mM Tris-HCl, 1 mM CaCl₂, 50% Glycerol, (pH 7.4 @ 25°C)
Specification Version: PS-P8107S v2.0

Proteinase K, Molecular Biology Grade Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
P8107SVIAL	Proteinase K, Molecular Biology Grade	10145029	Pass

Assay Name/Specification	Lot # 10156147
<p>qPCR DNA Contamination (Eukaryotic Genomic) A minimum of 1.6 units of Proteinase K, Molecular Biology Grade is screened for the presence of eukaryotic genomic DNA using SYBR® Green qPCR with universal primers for the 18S rRNA locus. Results are quantified using a standard curve generated from purified E. album genomic DNA. The measured level of eukaryotic genomic DNA contamination is ≤ 2.5 pg DNA/µl.</p>	Pass
<p>Single Stranded DNase Activity (FAM-Labeled Oligo) A 50 µl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent internal labeled oligonucleotide and a minimum of 4 units of Proteinase K, Molecular Biology Grade incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.</p>	Pass
<p>RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of fluorescein labeled RNA transcript and a minimum of 0.8 units of Proteinase K, Molecular Biology Grade is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass
<p>Non-Specific DNase Activity (16 Hour)</p>	Pass


Assay Name/Specification	Lot # 10156147
<p>A 50 µl reaction in CutSmart® Buffer containing 1 µg of Lambda-HindIII DNA and a minimum of 0.8 units of Proteinase K, Molecular Biology Grade incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	
<p>Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 8 units of Proteinase K, Molecular Biology Grade incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.</p>	Pass
<p>Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart® Buffer containing 1 µg of supercoiled PhiX174 RF I DNA and a minimum of 0.8 units of Proteinase K, Molecular Biology Grade incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Alicia Bielik
Production Scientist
23 Jun 2022



Erin Varney
Packaging Quality Control Inspector
23 Jun 2022