

## New England Biolabs Certificate of Analysis

**Product Name:** EcoRV  
**Catalog Number:** R0195M  
**Concentration:** 100,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 µl.  
**Packaging Lot Number:** 10130811  
**Expiration Date:** 08/2023  
**Storage Temperature:** -20°C  
**Storage Conditions:** 50 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA  
**Specification Version:** PS-R0195T/M v1.0

EcoRV Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0195MVIAl	EcoRV	10118280	Pass
B7024AVIAl	Gel Loading Dye, Purple (6X)	10126634	Pass
B6003SVIAl	NEBuffer™ r3.1	10116057	Pass

Assay Name/Specification	Lot # 10130811
<b>Blue-White Screening (Terminal Integrity)</b> A sample of LITMUS28i vector linearized with a 10-fold excess of EcoRV, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass
<b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in NEBuffer 3.1 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 20 units of EcoRV incubated for 4 hours at 37°C results in <50% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in NEBuffer 3.1 containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 100 units of EcoRV incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
<b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in NEBuffer 3.1 containing 1 µg of Lambda DNA and a minimum of 20 units of EcoRV incubated for 16 hours at 37°C results in a DNA pattern free of	Pass


Assay Name/Specification	Lot # 10130811
<p>detectable nuclease degradation as determined by agarose gel electrophoresis.</p> <p><b>Ligation and Recutting (Terminal Integrity)</b> After a 10-fold over-digestion of Lambda DNA with EcoRV, &gt;95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, &gt;95% can be recut with EcoRV.</p> <p><b>Protein Purity Assay (SDS-PAGE)</b> EcoRV is &gt;95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.</p>	<p style="text-align: center;"><b>Pass</b></p> <p style="text-align: center;"><b>Pass</b></p>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Penghua Zhang  
Production Scientist  
23 Nov 2021



Michael Tonello  
Packaging Quality Control Inspector  
23 Nov 2021