

## New England Biolabs Certificate of Analysis

**Product Name:** MseI  
**Catalog Number:** R0525S  
**Concentration:** 10,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 µl.  
**Lot Number:** 10031770  
**Expiration Date:** 10/2020  
**Storage Temperature:** -20°C  
**Storage Conditions:** 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA  
**Specification Version:** PS-R0525S/L v1.0

MseI Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0525SVIAL	MseI	10023121	Pass
B7204SVIAL	CutSmart® Buffer	10031557	Pass
B7024SVIAL	Gel Loading Dye, Purple (6X)	10021132	Pass

Assay Name/Specification	Lot # 10031770
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 30 units of MseI incubated for 4 hours at 37°C releases <0.2% of the total radioactivity.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> After a 10-fold over-digestion of Lambda DNA with MseI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with MseI.	Pass
<b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 100 Units of MseI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

*Stephanie Cornelio*

---

Stephanie Cornelio  
Production Scientist  
01 Oct 2018

*Michael Tonello*

---

Michael Tonello  
Packaging Quality Control Inspector  
19 Dec 2018