

## New England Biolabs Certificate of Analysis

**Product Name:** Bfal  
**Catalog Number:** R0568L  
**Concentration:** 10,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 µl.  
**Lot Number:** 10054781  
**Expiration Date:** 09/2021  
**Storage Temperature:** -80°C  
**Storage Conditions:** 300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 500 µg/ml BSA  
**Specification Version:** PS-R0568S/L v2.0


Bfal Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0568LVIAL	Bfal	10054782	Pass
B7204SVIAL	CutSmart® Buffer	10053981	Pass

Assay Name/Specification	Lot # 10054781
<p><b>Exonuclease Activity (Radioactivity Release)</b>            A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [<sup>3</sup>H] E. coli DNA and a minimum of 100 units of Bfal incubated for 4 hours at 37°C releases &lt;0.1% of the total radioactivity.</p>	Pass
<p><b>Ligation and Recutting (Terminal Integrity)</b>            After a 5-fold over-digestion of Lambda DNA with Bfal, ~25% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, &gt;95% can be recut with Bfal.</p>	Pass
<p><b>Non-Specific DNase Activity (16 Hour)</b>            A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 100 Units of Bfal incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.



Penghua Zhang  
Production Scientist  
10 Sep 2019



Michael Tonello  
Packaging Quality Control Inspector  
26 Sep 2019