

## New England Biolabs Product Specification

|                               |  |
|-------------------------------|--|
| <i>Product Name:</i>          | <i>Multiplex PCR 5X Master Mix</i>   |
| <i>Catalog #:</i>             | <i>M0284S</i>  |
| <i>Concentration:</i>         | <i>5X Concentrate</i>  |
| <i>Shelf Life:</i>            | <i>24 months</i>   |
| <i>Storage Temp:</i>          | <i>-20°C</i>   |
| <i>Composition (1X):</i>      | <i>20 mM Tris-HCl (pH 8.9 @ 25°C), 50 mM KCl, 30 mM NH<sub>4</sub>Cl, 2.5 mM MgCl<sub>2</sub>, 0.3 mM dATP, 0.3 mM dCTP, 0.3 mM dGTP, 0.3 mM dTTP, 3.2 % Glycerol, 0.08 % IGEPAL<sup>®</sup> CA-630, 0.07 % Tween<sup>®</sup> 20, 67 units/ml Taq DNA Polymerase</i> |
| <i>Specification Version:</i> | <i>PS-M0284S v1.0</i>  |
| <i>Effective Date:</i>        | <i>07 Jul 2016</i>   |

### Assay Name/Specification (minimum release criteria)

**Endonuclease Activity (Nicking)** - A 50 µl reaction in ThermoPol<sup>®</sup> Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 20 units of Taq DNA Polymerase incubated for 4 hours at either 37°C or 75°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.

**Non-Specific DNase Activity (16 hour, Buffer)** - A 50 µl reaction in 2X Multiplex PCR Master Mix containing 1 µg of T3 DNA in addition to a reaction containing Lambda-HindIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

**PCR Amplification (15-plex PCR, Master Mix)** - A 25 µl reaction in 1X Multiplex PCR Master Mix and 0.15 µM primer mix containing 10 ng Human Genomic DNA for 35 cycles of PCR amplification results in the expected 15 products.

**Phosphatase Activity (pNPP)** - A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl<sub>2</sub> containing 2.5 mM *p*-Nitrophenyl Phosphate (pNPP) and a minimum of 100 units of Taq DNA Polymerase incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.

**Protein Purity Assay (SDS-PAGE)** - Taq DNA Polymerase is ≥ 99% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.

**qPCR DNA Contamination (*E. coli* Genomic)** - A minimum of 5 units of Taq DNA Polymerase is screened for the presence of *E. coli* genomic DNA using SYBR<sup>®</sup> Green qPCR with primers specific for the *E. coli* 16S rRNA locus. Results are quantified using a standard curve generated from purified *E. coli* genomic DNA. The measured level of *E. coli* genomic DNA contamination is ≤ 1 *E. coli* genome.



## New England Biolabs Product Specification

**Assay Name/Specification (minimum release criteria)**

**RNase Activity (Extended Digestion)** - A 10  $\mu$ l reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1  $\mu$ l of Multiplex PCR 5X Master Mix is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.

**Single Stranded DNase Activity (FAM-Labeled Oligo)** - A 50  $\mu$ l reaction in ThermoPol<sup>®</sup> Reaction Buffer containing a 10 nM solution of a fluorescent internal labeled oligonucleotide and a minimum of 25 units of *Taq* DNA Polymerase incubated for 30 minutes at either 37°C or 75°C yields <10% degradation as determined by capillary electrophoresis.



Date 07 Jul 2016

Derek Robinson  
Director of Quality Control

