

## New England Biolabs Product Specification

<i>Product Name:</i>	<i>Hemo KlenTaq<sup>®</sup></i>
<i>Catalog #:</i>	<i>M0332S/L/V</i>
<i>Concentration:</i>	<i>500 reactions/ml</i>
<i>Unit Definition:</i>	<i>N/A</i>
<i>Shelf Life:</i>	<i>24 months</i>
<i>Storage Temp:</i>	<i>-20°C</i>
<i>Storage Conditions:</i>	<i>10 mM Tris-HCl, 100 mM KCl, 1 mM DTT, 0.1 mM EDTA, 0.5 % Tween<sup>®</sup> 20, 0.5 % IGEPAL<sup>®</sup> CA-630, 50 % Glycerol, (pH 7.4 @ 25°C)</i>
<i>Specification Version:</i>	<i>PS-M0332S/L v2.0</i>
<i>Effective Date:</i>	<i>12 Feb 2020</i>

### Assay Name/Specification (minimum release criteria)

**Endonuclease Activity (Nicking)** - A 50 µl reaction in Hemo KlenTaq<sup>®</sup> Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 8 µl of Hemo KlenTaq<sup>®</sup> incubated for 4 hours at 37°C and 75°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.

**Non-Specific DNase Activity (16 Hour)** - A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 1 µl of Hemo KlenTaq<sup>®</sup> incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

**PCR Amplification (0.5 kb Whole Blood DNA)** - A 50 µl reaction in Hemo KlenTaq<sup>®</sup> Reaction Buffer in the presence of 200 µM dNTPs and 0.3 µM primers containing 10% whole blood treated with sodium heparin, sodium EDTA, potassium EDTA or sodium citrate with 4 µl of Hemo KlenTaq<sup>®</sup> for 35 cycles of PCR amplification results in the expected 0.5 kb product.

**Phosphatase Activity (pNPP)** - A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl<sub>2</sub> containing 2.5 mM *p*-Nitrophenyl Phosphate (pNPP) and a minimum of 2 µl Hemo KlenTaq<sup>®</sup> incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.

**Protein Purity Assay (SDS-PAGE)** - Hemo KlenTaq<sup>®</sup> is ≥ 99% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.

**qPCR DNA Contamination (*E. coli* Genomic)** - A minimum of 1 µl of Hemo KlenTaq<sup>®</sup> is screened for the presence of *E. coli* genomic DNA using SYBR<sup>®</sup> Green qPCR with primers specific for the *E. coli* 16S rRNA locus. Results are quantified using a standard curve generated from purified *E. coli* genomic DNA. The measured level of *E. coli* genomic DNA contamination is ≤ 1 *E. coli* genome.



## New England Biolabs Product Specification

**Assay Name/Specification (minimum release criteria)**

**RNase Activity (Extended Digestion)** - A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Hemo KlenTaq<sup>®</sup> is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.

**Single Stranded DNase Activity (FAM-Labeled Oligo)** - A 20 µl reaction in Hemo KlenTaq<sup>®</sup> Reaction Buffer containing a 10 nM solution of a fluorescent internal labeled oligonucleotide and a minimum of 8 µl of Hemo KlenTaq<sup>®</sup> incubated for 30 minutes at 37°C and 75°C yields <10% degradation as determined by capillary electrophoresis.

*One or more products referenced in this document may be covered by a 3rd-party trademark.  
Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.*



Date 12 Feb 2020

Derek Robinson  
Director, Quality Control

